

Stark County's Influenza Weekly Report

Week 19 (5/8/22-5/14/2022)




Public Health
Prevent. Promote. Protect.

During Week 19:

- Stark County had 6 influenza-associated hospitalizations.
- Nationally, 0 influenza-associated pediatric deaths were reported.
- During Week 19, the following specimens were collected nationally from public health laboratories:

Type/Subtype	Week 19	Week 19 %
Influenza A	433	99.3%
(H1N1) pdm09	1	0.4%
H3N2	255	99.6%
H3N2v	0	0.0%
The remainder were not subtyped		
Influenza B	3	0.7%
Yamagata	0	0.0%
Victoria	0	0.0%
The remainder were not subtyped		

% of Specimens by Type Among Stark County Reported Influenza Cases




■ Influenza A % ■ Influenza B %

So Far this Year:

- There have been 115 influenza-associated hospitalizations in Stark County during this influenza season.
 - Hospitalizations range from 3 weeks to 100 years old with the median age being 67 years old.
- The national cumulative rate of influenza-associated hospitalizations is 14.4 per 100,000.
- Nationally, 24 influenza-associated pediatric deaths have been reported so far this year.
- Up to Week 19, the following specimens were collected nationally from public health laboratories:



Type/Subtype	YTD	YTD %
Influenza A	21,134	99.4%
(H1N1) pdm09	18	0.1%
H3N2	16,362	99.9%
H3N2v	1	<0.1%
The remainder were not subtyped		
Influenza B	120	0.6%
Yamagata	1	2.6%
Victoria	38	97.4%
The remainder were not subtyped		

DISTRIBUTION OF INFLUENZA-ASSOCIATED HOSPITALIZATIONS BY AGE GROUP WITH AGE-ADJUSTED RATE IN STARK COUNTY




Graph 1A and 1B: Hospitalized and Non-Hospitalized Influenza Cases Reported to the Local Health Department.

There were 20 cases of hospitalized and non-hospitalized influenza reported for Week 19 in Stark County. Last year, there were 0 cases reported during this week. Stark County has reported 595 cases so far this year compared to the 6 cases reported by this time last year.



Graph 2: Sentinel Provider Reported Influenza Like Illness (ILI) in Stark County.


- ◆ During Week 19, Stark County Sentinel Providers reported that 0.72% of office visits were due to ILI. The national reports show that 2.4% of outpatient visits were due to ILI, which is slightly below the national baseline of 2.5%.



Graph 3: Emergency Department Visits for Respiratory and Constitutional Syndromes.

(Source: Health Monitoring Systems, EpiCenter, Hospital and Stat Care Patient Registration Surveillance System)


- ◆ ED and Stat Care visits for respiratory and constitutional syndrome were reported at 24.86% of all visits, which was above the multi-year baseline average during Week 19.



Graph 4: Emergency Department (ED) Visits for Influenza Like Illness (+Fever).

(Source: Health Monitoring Systems, EpiCenter, Hospital and Stat Care Patient Registration Surveillance System)

- ◆ ED and Stat Care visits for ILI + Fever were reported at 2.69% of all visits, which was above the multi-year baseline average during Week 19.




- All data presented in this weekly report are preliminary and may change as additional information is received.
- Multiyear averages included in this weekly report do not include the 2009/2010 H1N1 season.
- If you have any questions please contact Kaelyn Boyd at 234.458.5135 or kboyd@cantonhealth.org or Chelsea Sadinski at 330.489.9914 or sadinskic@starkhealth.org

Graph 5: Stark County School Absenteeism.

(Source: Voluntary reporting by multiple school districts throughout the county)

- ◆ During Week 19, Stark County schools reported a total median absenteeism of 5.65%. Absenteeism due to influenza-like-illness was reported at 0.00% of all absences.




Map: ILI Activity Level Indicator Determined by Data Reported to ILINet.

(Data collected in ILINet may disproportionately represent certain populations within a jurisdiction or CBSA, and therefore, may not accurately depict the full picture of influenza activity for the entire jurisdiction or CBSA. Differences in the data presented here by CDC and independently by some health departments likely represent differing levels of data completeness with data presented by the health department likely being the more complete.)

A Weekly Influenza Surveillance Report Prepared by the Influenza Division Influenza-Like Illness (ILI) Activity Level Indicator Determined by Data Reported to ILINet

This system monitors visits for ILI (fever and cough or sore throat), not laboratory confirmed influenza and may capture patient visits due to other respiratory pathogens that cause similar symptoms.

2021-2022 Influenza Season Week 19 ending May 14, 2022



Viruses circulating vs. Viruses in 2021-2022 Influenza Season Vaccine:

According to an MMWR Report published on March 11th, 2022, this year's influenza vaccine is 16% effective at preventing laboratory-confirmed, mild/moderate (outpatient) medically attended influenza (95% Confidence Interval [CI]= -16% to 39%). This is from data collected on 3,636 children and adults enrolled in the U.S Influenza Vaccine Effectiveness Network (U.S Flu VE Network) during October 4, 2021- February 12, 2022. The CDC routinely recommends that health care providers continue to administer influenza vaccines to persons ages 6 months and older as long as influenza viruses are circulating, as the vaccine can prevent serious outcomes (<https://www.cdc.gov/mmwr/volumes/71/wr/mm7110a1.htm>).



- All data presented in this weekly report are preliminary and may change as additional information is received.
- Multiyear averages included in this weekly report do not include the 2009/2010 H1N1 season.
- If you have any questions please contact Kaelyn Boyd at 234.458.5135 or kboyd@cantonhealth.org or Chelsea Sadinski at 330.489.9914 or sadinskic@starkhealth.org